

# ASYMPTOTICALLY NONEXPANSIVE MAPPINGS IN MODULAR FUNCTION SPACES

T. DOMINGUEZ-BENAVIDES, M.A. KHAMSI AND S. SAMADI

## ABSTRACT

In this paper, we prove that if  $\rho$  is a convex,  $\sigma$ -finite modular function satisfying a  $\Delta_2$ -type condition,  $C$  a convex,  $\rho$ -bounded,  $\rho$ -a.e. compact subset of  $L_\rho$  and  $T : C \rightarrow C$  a  $\rho$ -asymptotically nonexpansive mapping, then  $T$  has a fixed point. In particular, any asymptotically nonexpansive self-map defined on a convex subset of  $L^1(\Omega, \mu)$  which is compact for the topology of convergence local in measure has a fixed point.

**1991 Mathematics subject classification** : Primary 46E30; Secondary 47H09, 47H10.

**Key Words:** asymptotically nonexpansive mappings, fixed point, modular functions.

---

The first author is partially supported by PB-96-1338-C01-C02 and PAI-FMQ-0127.

## INTRODUCTION

Let  $(M, d)$  a metric space. A mapping,  $T : M \rightarrow M$  is said to be asymptotically nonexpansive if there exists a sequence  $\{k_n\}$  of real numbers with  $\lim_{n \rightarrow \infty} k_n = 1$  such that

$$d(T^n x, T^n y) \leq k_n d(x, y)$$

for any  $x, y \in M$  and  $n \in \mathbb{N}$ . In 1970 Goebel and Kirk [5] proved that  $T$  has a fixed point whenever  $M$  is a convex bounded closed subset of a Banach space  $X$ . Further generalizations of this result were proved by Yu and Dai [14] when  $X$  is 2-uniformly rotund, by Martínez Yañez [10] and Xu [12] when  $X$  is  $k$ -uniformly rotund for some  $k \geq 1$ , by Xu [13] when  $X$  is nearly uniformly convex and by Kim and Xu [9] when  $X$  has uniform normal structure. Some special studies on the theory of the fixed point for asymptotically nonexpansive mappings were made by many other authors (see, for example, [2,11]).

The first fixed point results in modular function spaces were given by Khamsi, Kozłowski and Reich [7]. Even though a metric is not defined, many problems in metric fixed point theory can be reformulated in modular spaces. For instance, fixed point theorems are proved in [6,7] for nonexpansive mappings, in [3] for asymptotically regular mappings and in [4] for uniformly Lipschitzian mappings. In this paper we will prove the existence of fixed points for asymptotically nonexpansive mappings in modular function spaces when the modular  $\rho$  satisfies some convexity and  $\Delta_2$ -type properties.

Our results can be, in particular, applied to  $L^1(\Omega, \mu)$ , showing that asymptotically nonexpansive mappings have a fixed point when they are defined on a convex subset of  $L^1(\Omega, \mu)$  which is compact with respect to the topology of convergence local in measure.

## 1. PRELIMINARIES

We start by reviewing some basic facts about modular spaces as formulated by Kozłowski [8]. For more details the reader may consult [6,7].

Let  $\Omega$  be a nonempty set and  $\Sigma$  be a nontrivial  $\sigma$ -algebra of subsets of  $\Omega$ . Let  $\mathcal{P}$  be a  $\delta$ -ring of subsets of  $\Sigma$ , such that  $E \cap A \in \mathcal{P}$  for any  $E \in \mathcal{P}$  and  $A \in \Sigma$ . Let us assume that there exists an increasing sequence of sets  $K_n \in \mathcal{P}$  such that

$\Omega = \bigcup K_n$ . By  $\mathcal{E}$  we denote the linear space of all simple functions with supports from  $\mathcal{P}$ . By  $\mathcal{M}$  we will denote the space of all measurable functions, i.e. all functions  $f : \Omega \rightarrow \mathbb{R}$  such that there exists a sequence  $\{g_n\} \in \mathcal{E}$ ,  $|g_n| \leq |f|$  and  $g_n(\omega) \rightarrow f(\omega)$  for all  $\omega \in \Omega$ . By  $1_A$  we denote the characteristic function of the set  $A$ .

**Definition 1.1.** A functional  $\rho : \mathcal{E} \times \Sigma \rightarrow [0, \infty]$  is called a function modular if:

- (P<sub>1</sub>)  $\rho(0, E) = 0$  for any  $E \in \Sigma$ ,
- (P<sub>2</sub>)  $\rho(f, E) \leq \rho(g, E)$  whenever  $|f(\omega)| \leq |g(\omega)|$  for any  $\omega \in \Omega$ ,  $f, g \in \mathcal{E}$  and  $E \in \Sigma$ ,
- (P<sub>3</sub>)  $\rho(f, \cdot) : \Sigma \rightarrow [0, \infty]$  is a  $\sigma$ -subadditive measure for every  $f \in \mathcal{E}$ ,
- (P<sub>4</sub>)  $\rho(\alpha, A) \rightarrow 0$  as  $\alpha$  decreases to 0 for every  $A \in \mathcal{P}$ , where  $\rho(\alpha, A) = \rho(\alpha 1_A, A)$ ,
- (P<sub>5</sub>) if there exists  $\alpha > 0$  such that  $\rho(\alpha, A) = 0$ , then  $\rho(\beta, A) = 0$  for every  $\beta > 0$ ,
- (P<sub>6</sub>) for any  $\alpha > 0$   $\rho(\alpha, \cdot)$  is order continuous on  $\mathcal{P}$ , that is  $\rho(\alpha, A_n) \rightarrow 0$  if  $\{A_n\} \in \mathcal{P}$  and decreases to  $\emptyset$ .

The definition of  $\rho$  is then extended to  $f \in \mathcal{M}$  by

$$\rho(f, E) = \sup\{\rho(g, E); g \in \mathcal{E}, |g(\omega)| \leq |f(\omega)| \text{ for every } \omega \in \Omega\}.$$

**Definition 1.2.** A set  $E$  is said to be  $\rho$ -null if  $\rho(\alpha, E) = 0$  for every  $\alpha > 0$ .

For the sake of simplicity we write  $\rho(f)$  instead of  $\rho(f, \Omega)$ .

**Definition 1.3.** A modular function  $\rho$  is called  $\sigma$ -finite if there exists an increasing sequence of sets  $K_n \in \mathcal{P}$  such that  $0 < \rho(K_n) < \infty$  and  $\Omega = \bigcup K_n$ .

It is easy to see that the functional  $\rho : \mathcal{M} \rightarrow [0, \infty]$  is a modular and satisfies the following properties:

- (i)  $\rho(f) = 0$  iff  $f = 0$   $\rho$ -a.e.
- (ii)  $\rho(\alpha f) = \rho(f)$  for every scalar  $\alpha$  with  $|\alpha| = 1$  and  $f \in \mathcal{M}$ .
- (iii)  $\rho(\alpha f + \beta g) \leq \rho(f) + \rho(g)$  if  $\alpha + \beta = 1$ ,  $\alpha \geq 0, \beta \geq 0$  and  $f, g \in \mathcal{M}$ .

In addition, if the following property is satisfied

(iii)'  $\rho(\alpha f + \beta g) \leq \alpha\rho(f) + \beta\rho(g)$  if  $\alpha + \beta = 1$  ;  $\alpha \geq 0, \beta \geq 0$  and  $f, g \in \mathcal{M}$ ,

we say that  $\rho$  is a convex modular.

The modular  $\rho$  defines a corresponding modular space, i.e the vector space  $L_\rho$  given by

$$L_\rho = \{f \in \mathcal{M}; \rho(\lambda f) \rightarrow 0 \text{ as } \lambda \rightarrow 0\}.$$

When  $\rho$  is convex, the formula

$$\|f\|_\rho = \inf \left\{ \alpha > 0; \rho\left(\frac{f}{\alpha}\right) \leq 1 \right\}$$

defines a norm in the modular space  $L_\rho$  which is frequently called the Luxemburg norm. We can also consider the space

$$E_\rho = \{f \in \mathcal{M}; \rho(\alpha f, A_n) \rightarrow 0 \text{ as } n \rightarrow \infty \text{ for every } A_n \in \Sigma \text{ that decreases to } \emptyset \text{ and } \alpha > 0\}.$$

**Definition 1.4.** A function modular is said to satisfy the  $\Delta_2$ -condition if

$$\sup_{n \geq 1} \rho(2f_n, D_k) \rightarrow 0 \text{ as } k \rightarrow \infty \text{ whenever } \{f_n\}_{n \geq 1} \subset \mathcal{M}, D_k \in \Sigma \text{ decreases to } \emptyset \text{ and } \sup_{n \geq 1} \rho(f_n, D_k) \rightarrow 0 \text{ as } k \rightarrow \infty.$$

We know from [8] that  $E_\rho = L_\rho$  when  $\rho$  satisfies the  $\Delta_2$ -condition.

**Definition 1.5.** A function modular is said to satisfy the  $\Delta_2$ -type condition if there exists  $K > 0$  such that for any  $f \in L_\rho$  we have  $\rho(2f) \leq K\rho(f)$ .

In general,  $\Delta_2$ -type condition and  $\Delta_2$ -condition are not equivalent, even though it is obvious that  $\Delta_2$ -type condition implies  $\Delta_2$ -condition on the modular space  $L_\rho$ .

**Definition 1.6.** Let  $L_\rho$  be a modular space.

- (1) The sequence  $\{f_n\}_n \subset L_\rho$  is said to be  $\rho$ -convergent to  $f \in L_\rho$  if  $\rho(f_n - f) \rightarrow 0$  as  $n \rightarrow \infty$ .
- (2) The sequence  $\{f_n\}_n \subset L_\rho$  is said to be  $\rho$ -a.e. convergent to  $f \in L_\rho$  if the set  $\{\omega \in \Omega; f_n(\omega) \not\rightarrow f(\omega)\}$  is  $\rho$ -null.

- (3) The sequence  $\{f_n\}_n \subset L_\rho$  is said to be  $\rho$ -Cauchy if  $\rho(f_n - f_m) \rightarrow 0$  as  $n$  and  $m$  go to  $\infty$ .
- (4) A subset  $C$  of  $L_\rho$  is called  $\rho$ -closed if the  $\rho$ -limit of a  $\rho$ -convergent sequence of  $C$  always belongs to  $C$ .
- (5) A subset  $C$  of  $L_\rho$  is called  $\rho$ -a.e. closed if the  $\rho$ -a.e. limit of a  $\rho$ -a.e. convergent sequence of  $C$  always belongs to  $C$ .
- (6) A subset  $C$  of  $L_\rho$  is called  $\rho$ -a.e. compact if every sequence in  $C$  has a  $\rho$ -a.e. convergent subsequence in  $C$ .
- (7) A subset  $C$  of  $L_\rho$  is called  $\rho$ -bounded if

$$\delta_\rho(C) = \sup\{\rho(f - g); f, g \in C\} < \infty.$$

We recall two basic results (see [7]) in the theory of modular spaces.

- (i) If there exists a number  $\alpha > 0$  such that  $\rho(\alpha(f_n - f)) \rightarrow 0$ , then there exists a subsequence  $\{g_n\}_n$  of  $\{f_n\}_n$  such that  $g_n \rightarrow f$   $\rho$ -a.e.
- (ii) (Lebesgue's Theorem) If  $f_n, f \in \mathcal{M}$ ,  $f_n \rightarrow f$   $\rho$ -a.e. and there exists a function  $g \in E_\rho$  such that  $|f_n| \leq |g|$   $\rho$ -a.e. for all  $n$ , then  $\|f_n - f\|_\rho \rightarrow 0$ .

We know, by [6,7] that under  $\Delta_2$ -condition the norm convergence and modular convergence are equivalent, which implies that the norm and modular convergence are also the same when we deal with the  $\Delta_2$ -type condition.

In the sequel we will assume that the modular function  $\rho$  is convex and satisfies the  $\Delta_2$ -type condition.

**Definition 1.7.** Let  $\rho$  be as above. We define a growth function  $\omega$  by:

$$\omega(t) = \sup \left\{ \frac{\rho(tf)}{\rho(f)}, f \in L_\rho \setminus \{0\} \right\} \quad \text{for all } 0 \leq t < \infty.$$

We have the following:

**Lemma 1.1.** [3] *Let  $\rho$  be as above. Then the growth function  $\omega$  has the following properties:*

- (1)  $\omega(t) < \infty, \forall t \in [0, \infty)$

(2)  $\omega : [0, \infty) \rightarrow [0, \infty)$  is a convex, strictly increasing function. So, it is continuous.

(3)  $\omega(\alpha\beta) \leq \omega(\alpha)\omega(\beta); \forall \alpha, \beta \in [0, \infty)$

(4)  $\omega^{-1}(\alpha)\omega^{-1}(\beta) \leq \omega^{-1}(\alpha\beta); \forall \alpha, \beta \in [0, \infty)$ , where  $\omega^{-1}$  is the function inverse of  $\omega$ .

The following lemma shows that the growth function can be used to give an upper bound for the norm of a function.

**Lemma 1.2.** [3] *Let  $\rho$  be a convex function modular satisfying the  $\Delta_2$ -type condition. Then*

$$\|f\|_\rho \leq \frac{1}{\omega^{-1}\left(\frac{1}{\rho(f)}\right)} \quad \text{whenever } f \in L_\rho.$$

The next lemma will be of major interest throughout this work.

**Lemma 1.3.** [6] *Let  $\rho$  be a function modular satisfying the  $\Delta_2$ -condition and  $\{f_n\}_n$  be a sequence in  $L_\rho$  such that  $f_n \xrightarrow{\rho\text{-a.e.}} f \in L_\rho$  and there exists  $k > 1$  such that  $\sup_n \rho(k(f_n - f)) < \infty$ . Then,*

$$\liminf_{n \rightarrow \infty} \rho(f_n - g) = \liminf_{n \rightarrow \infty} \rho(f_n - f) + \rho(f - g) \quad \text{for all } g \in L_\rho.$$

Moreover, we have

$$\rho(f) \leq \liminf_{n \rightarrow \infty} \rho(f_n).$$

## 2. AN EQUIVALENT TOPOLOGY

The concept of  $\rho$ -a.e. closed, compact sets have been studied extensively in the sequential case. One of the problem that many authors have found hard to circumvent is whether these notions are related to a topology. In this section we will discuss this problem. In particular, we will construct a topology  $\tau$  for which the  $\rho$ -a.e. compactness is equivalent to the usual compactness for  $\tau$ . This is crucial when we try to use Zorn's lemma.

From now on, we assume that the modular function  $\rho$  is, in addition,  $\sigma$ -finite. Set

$$d(f, g) = \sum_{k=1}^{\infty} \frac{1}{2^k} \frac{1}{\rho(1_{K_k})} \rho \left( \frac{|f - g|}{1 + |f - g|} 1_{K_k} \right) \quad \text{for any } f, g \in L_\rho.$$

Some basic properties satisfied by  $d$  are discussed in the following proposition.

**Proposition 2.1.** The functional  $d$  satisfies the following:

- (1)  $d(f, g) = 0$  if and only if  $f = g$   $\rho$ -a.e.;
- (2)  $d(f, g) = d(g, f)$ ;
- (3)  $d(f, g) \leq \frac{\omega(2)}{2} (d(f, h) + d(h, g))$ ;

for any  $f, g$  and  $h$  in  $L_\rho$ .

*Proof.* (1) and (2) are obvious. To prove (3) we only need to recall the inequality

$$\frac{|a + b|}{1 + |a + b|} \leq \frac{|a|}{1 + |a|} + \frac{|b|}{1 + |b|}$$

for all positive numbers  $a, b$  and use the definition of the growth function  $\omega$ . □

**Remark 2.1.** The functional  $d$  is not a distance because of (3). But there are many mathematical objects which fails the triangle inequality but are very useful tools. That is the case with  $d$ .

In the next proposition, we discuss the relationship between  $\rho$ -a.e. convergence and the convergence for the functional  $d$ .

**Proposition 2.2.** Let  $\rho$  be a convex,  $\sigma$ -finite modular satisfying the  $\Delta_2$ -type condition and  $\{f_n\}_n$  be a sequence of measurable functions. If  $\{f_n\}_n$  is  $\rho$ -a.e. convergent to  $f$ , then

$$\lim_{n \rightarrow \infty} d(f_n, f) = 0.$$

Moreover, if

$$\lim_{n \rightarrow \infty} d(f_n, f) = 0,$$

then there exists a subsequence  $\{f_{n_k}\}_k$  which converges  $\rho$ -a.e. to  $f$ .

*Proof.* Assume that  $\{f_n\}_n$   $\rho$ -a.e. converges to  $f$ . We will show that  $\lim_{n \rightarrow \infty} d(f_n, f) =$

0. Let  $\varepsilon > 0$ , and choose  $N \in \mathbb{N}$  such that  $\sum_{k=N+1}^{\infty} \frac{1}{2^k} < \varepsilon$ . We have

$$\begin{aligned} \lim_{n \rightarrow \infty} d(f_n, f) &\leq \lim_{n \rightarrow \infty} \sum_{k=1}^N \frac{1}{2^k} \frac{1}{\rho(1_{K_k})} \rho \left( \frac{|f_n - f|}{1 + |f_n - f|} 1_{K_k} \right) + \varepsilon \\ &= \sum_{k=1}^N \lim_{n \rightarrow \infty} \frac{1}{2^k} \frac{1}{\rho(1_{K_k})} \rho \left( \frac{|f_n - f|}{1 + |f_n - f|} 1_{K_k} \right) + \varepsilon. \end{aligned}$$

Since

$$\frac{|f_n - f|}{1 + |f_n - f|} 1_{K_k} \xrightarrow{\rho\text{-a.e.}} 0 \quad \text{as } n \rightarrow \infty$$

for any  $k \in \mathbb{N}$  and  $\frac{|f_n - f|}{1 + |f_n - f|} 1_{K_k} \leq 1_{K_k}$ , from Lebesgue's Theorem we obtain

$$\lim_{n \rightarrow \infty} \rho \left( \frac{|f_n - f|}{1 + |f_n - f|} 1_{K_k} \right) = 0 \text{ for every non null integer } k. \text{ Thus } \lim_{n \rightarrow \infty} d(f_n, f) \leq \varepsilon$$

for each  $\varepsilon > 0$  which means that  $\lim_{n \rightarrow \infty} d(f_n, f) = 0$ .

Assume now that  $\lim_{n \rightarrow \infty} d(f_n, f) = 0$ . For every non null integer  $k$  we have

$$\lim_{n \rightarrow \infty} \rho \left( \frac{|f_n - f|}{1 + |f_n - f|} 1_{K_k} \right) = 0.$$

Thus, there exists a subsequence  $\{f_n^1\}_n$  of  $\{f_n\}_n$  such that  $\frac{|f_n^1 - f|}{1 + |f_n^1 - f|} 1_{K_1} \xrightarrow{\rho\text{-a.e.}} 0$

and so  $f_n^1 \xrightarrow{\rho\text{-a.e.}} f$  in  $K_1$  i.e.  $\lim_{n \rightarrow \infty} f_n^1(x) = f(x)$  whenever  $x \in K_1 \setminus A_1$  where  $A_1 \subset K_1$  and  $\rho(1_{A_1}) = 0$ .

By induction and using a diagonal argument we obtain a subsequence of  $\{f_n\}_n$  which converges  $\rho$ -a.e. to  $f$ .  $\square$

**Definition 2.1.** Let  $C$  be a subset of  $L_\rho$ .

- (a)  $C$  is said to be  $d$ -closed iff for any sequence  $\{f_n\}_n$  in  $C$  which  $d$ -converges to  $f$ , then we have  $f \in C$ .
- (b)  $C$  is  $d$ -open iff  $L_\rho \setminus C$  is  $d$ -closed.

- (c)  $C$  is said to be  $d$ -sequentially compact if for each sequence  $\{f_n\}_n$  there exists a subsequence  $\{f_{n_k}\}_k$  which  $d$ -converges to a point in  $C$ .

It is easily seen that the family of all  $d$ -open subsets of  $L_\rho$  form a topology on  $L_\rho$ . Furthermore, from proposition (2.2)  $d$ -sequentially compact sets and  $\rho$ -a.e. compact sets are identical. On the other hand, even though  $d$  satisfies (3) instead of the triangular inequality, the usual arguments which prove that sequential compactness and compactness are identical in metric spaces hold in this setting. We also have  $d$ -sequential compactness and  $d$ -compactness are identical.

### 3. TECHNICAL LEMMAS

In the sequel we assume that  $\rho$  is a convex,  $\sigma$ -finite modular function satisfying the  $\Delta_2$ -type condition,  $C$  is a convex,  $\rho$ -bounded and  $\rho$ -a.e. compact subset of the modular function space  $L_\rho$  and  $T : C \rightarrow C$  is a  $\rho$ -asymptotically nonexpansive mapping, i.e. there exists a sequence of positive integers  $\{k_n\}_n$  which converge to 1 such that for every  $n \in \mathbb{N}$  and  $f, g \in C$  we have  $\rho(T^n f - T^n g) \leq k_n \rho(f - g)$ .

**Lemma 3.1.** *Under the above assumptions, let  $\{f_n\}_n$  be a sequence of elements of  $C$ . Consider the functional  $\Phi : C \rightarrow R$  defined by  $\Phi(g) = \limsup_{n \rightarrow \infty} \rho(f_n - g)$ .*

*Then, for any sequence  $\{g_m\}_m$  in  $C$  which  $\rho$ -a.e. converges to  $g \in C$  we have*

$$\Phi(g) \leq \liminf_{m \rightarrow \infty} \Phi(g_m).$$

*Proof.* Since  $C$  is  $\rho$ -a.e. compact, there exists a subsequence  $\{f_{\phi(n)}\}_n$  of  $\{f_n\}_n$  such that  $f_{\phi(n)} \xrightarrow{\rho\text{-a.e.}} f \in C$  and  $\lim_{n \rightarrow \infty} \rho(f_{\phi(n)} - g) = \limsup_{n \rightarrow \infty} \rho(f_n - g)$ . Hence

$$\begin{aligned} \Phi(g_m) &= \limsup_{n \rightarrow \infty} \rho(f_n - g_m) \\ &\geq \limsup_{n \rightarrow \infty} \rho(f_{\phi(n)} - g_m) \\ &\geq \liminf_{n \rightarrow \infty} \rho(f_{\phi(n)} - g_m). \end{aligned}$$

Lemma (1.3) implies

$$\liminf_{n \rightarrow \infty} \rho(f_{\phi(n)} - g_m) = \liminf_{n \rightarrow \infty} \rho(f_{\phi(n)} - f) + \rho(f - g_m).$$

Thus,  $\Phi(g_m) \geq \liminf_{n \rightarrow \infty} \rho(f_{\phi(n)} - f) + \rho(f - g_m)$ , for any  $m \leq 1$ . Hence

$$\liminf_{m \rightarrow \infty} \Phi(g_m) \geq \liminf_{n \rightarrow \infty} \rho(f_{\phi(n)} - f) + \liminf_{m \rightarrow \infty} \rho(f - g_m).$$

Again using lemma (1.3), we have

$$\liminf_{m \rightarrow \infty} \rho(f - g_m) = \liminf_{m \rightarrow \infty} \rho(g_m - g) + \rho(g - f),$$

which implies

$$\liminf_{m \rightarrow \infty} \Phi(g_m) \geq \liminf_{n \rightarrow \infty} \rho(f_{\phi(n)} - f) + \liminf_{m \rightarrow \infty} \rho(g_m - g) + \rho(g - f) \quad (I).$$

On the other hand,

$$\Phi(g) = \limsup_{n \rightarrow \infty} \rho(f_n - g) = \lim_{n \rightarrow \infty} \rho(f_{\phi(n)} - g) = \liminf_{n \rightarrow \infty} \rho(f_{\phi(n)} - g)$$

which implies

$$\Phi(g) = \liminf_{n \rightarrow \infty} \rho(f_{\phi(n)} - f) + \rho(f - g) \quad (II).$$

From (I) and (II), it is clear that

$$\Phi(g) \leq \liminf_{m \rightarrow \infty} \Phi(g_m),$$

which completes the proof. □

Denote  $\mathfrak{S}$  the family of all subsets  $K$  of  $C$  satisfying the following property:  $K$  is a nonempty, convex and  $\rho$ -a.e. closed subset of  $C$  such that

$$f \in K \quad \text{implies} \quad \Omega_{\rho\text{-a.e.}}(f) \subset K \quad (3.1)$$

where  $\Omega_{\rho\text{-a.e.}}(f) = \{g \in L_\rho : g = \lim_{i \rightarrow \infty} T^{n_i}(f) \text{ } \rho\text{-a.e for some } n_i \uparrow \infty\}$ . Ordering  $\mathfrak{S}$  by inclusion, there exists a nonempty minimal element  $H$  in  $\mathfrak{S}$  which satisfies (3.1) by using Zorn's lemma because  $C$  is compact for the topology generated by  $d$ .

The following lemma is the counterpart in modular function spaces of lemma (2.1) in [13] for Banach spaces.

**Lemma 3.2.** *Under the above assumptions, for each  $f \in H$  define the functional*

$$r_f(g) = \limsup_{n \rightarrow \infty} \rho(T^n f - g)$$

for any  $g \in L_\rho$ . Then the functional  $r_f(\cdot)$  is constant on  $H$  and this constant is independent of  $f$  in  $H$ .

*Proof.* Let  $t > 0$  and  $f \in H$ . Set

$$H_t(f) = \{ g \in H, \quad r_f(g) \leq t \}.$$

It is easily seen that  $H_t(f)$  is convex. We claim that  $H_t(f)$  is  $\rho$ -a.e. closed. Indeed, assume that  $\{g_m\}_m \in H_t(f)$   $\rho$ -a.e. converges to  $g \in H$ . Using Lemma (3.1), we get

$$\limsup_{n \rightarrow \infty} \rho(T^n f - g) \leq \liminf_{m \rightarrow \infty} \limsup_{n \rightarrow \infty} \rho(T^n f - g_m) \leq t.$$

Hence  $g \in H_t(f)$ , which clearly implies that  $H_t(f)$  is  $\rho$ -a.e. closed. Since  $H$  is  $\rho$ -a.e. compact we have that  $H_t(f)$  is  $\rho$ -a.e. compact. Next, we claim that  $H_t(f)$  satisfies property (3.1). Indeed, let  $g \in H_t(f)$  and  $h \in \Omega_{\rho\text{-a.e.}}(g)$ . We need to check that  $h \in H_t(f)$ . By definition of  $\Omega_{\rho\text{-a.e.}}(g)$ , there exists an increasing sequence of integers  $\{n_i\}_i$  such that  $T^{n_i}(g) \xrightarrow{\rho\text{-a.e.}} h$ . Lemma (3.1) implies

$$\begin{aligned} r_f(h) &= \limsup_{n \rightarrow \infty} \rho(T^n f - h) \leq \liminf_{i \rightarrow \infty} \limsup_{n \rightarrow \infty} \rho(T^n f - T^{n_i} g) \\ &\leq \liminf_{i \rightarrow \infty} r_f(T^{n_i}(g)) \leq \limsup_{i \rightarrow \infty} r_f(T^{n_i}(g)) \leq \limsup_{m \rightarrow \infty} r_f(T^m(g)) \\ &\leq \limsup_{m \rightarrow \infty} \left( \limsup_{n \rightarrow \infty} \rho(T^n f - T^m g) \right) \\ &\leq \limsup_{m \rightarrow \infty} \left( k_m \limsup_{n \rightarrow \infty} \rho(T^{n-m} f - g) \right) \\ &\leq \limsup_{m \rightarrow \infty} \limsup_{n \rightarrow \infty} \rho(T^n f - g) \leq t. \end{aligned}$$

Hence  $h \in H_t(f)$  as claimed. The minimality of  $H$  implies that  $H_t(f)$  is  $\emptyset$  or equal to  $H$ . From this, it is clear that  $r_t(\cdot)$  is constant on  $H$ . In order to complete the proof of this lemma, we need to prove that  $r_f$  is independent of  $f$ . Let  $f, g \in H$ . Since  $C$  is  $\rho$ -a.e. compact, there exists a subsequence  $\{T^{n_i}(g)\}_i$  of  $\{T^n(g)\}_n$  which  $\rho$ -a.e. converges to  $h \in C$ . Since  $H$  satisfies property (3.1), we have  $h \in H$ . Lemma (1.3) implies

$$\rho(T^n f - h) \leq \liminf_{i \rightarrow \infty} \rho(T^n f - T^{n_i} g).$$

Hence

$$\begin{aligned}
r_f &= r_f(h) = \limsup_{n \rightarrow \infty} \rho(T^n f - h) \\
&\leq \limsup_{n \rightarrow \infty} \liminf_{i \rightarrow \infty} \rho(T^n f - T^{ni} g) \\
&\leq \limsup_{n \rightarrow \infty} \limsup_{m \rightarrow \infty} \rho(T^n f - T^m g) \\
&\leq \limsup_{m \rightarrow \infty} \rho(f - T^m g) = r_g(f) = r_g,
\end{aligned}$$

which obviously implies  $r_g = r_f$ .  $\square$

Recall that if  $\rho$  satisfies the  $\Delta_2$ -type condition, then  $\rho$ -convergence and norm (i.e. Luxemburg norm) convergence coincide. We have the following result:

**Lemma 3.3.** *Let  $\rho$  be a convex modular function satisfying the  $\Delta_2$ -type condition. Let  $S$  be a nonempty, norm-compact subset of  $L_\rho$  with  $\text{diam}_\rho(S) > 0$ . Then there exists  $f \in \overline{\text{conv}}(S)$  such that*

$$\sup\{\rho(g - f) : g \in S\} < \text{diam}_\rho(S).$$

*Proof.* The proof is similar to the classical one known in Banach spaces. Indeed, since  $S$  is compact and  $\rho$  is norm continuous, there exist  $f_0, f_1 \in S$  such that  $\rho(f_0 - f_1) = \text{diam}_\rho(S)$ . Let  $S_0$  be a maximal subset of  $S$  such that  $f_0, f_1 \in S_0$  and for any  $f, g \in S_0$ ,  $f \neq g$ , we have  $\rho(f - g) = \text{diam}_\rho(S)$ . Since  $S$  is compact,  $S_0$  must be finite. Write  $S_0 = \{f_0, f_1, f_2, \dots, f_n\}$  and define

$$h = \frac{f_0 + f_1 + \dots + f_n}{n + 1}.$$

Since  $S$  is compact, there exists  $g_0 \in S$  such that

$$\rho(g_0 - h) = \sup\{\rho(g - h) : g \in S\}.$$

On the other hand, using the convexity of  $\rho$ , we get

$$\begin{aligned}
\rho(g_0 - h) &= \rho\left(\sum_{k=0}^{k=n} \left(\frac{1}{n+1}\right) g_0 - \sum_{k=0}^{k=n} \left(\frac{1}{n+1}\right) f_k\right) \\
&\leq \sum_{k=0}^{k=n} \left(\frac{1}{n+1}\right) \rho(g_0 - f_k) \leq \text{diam}_\rho(S).
\end{aligned}$$

If  $\rho(g_0 - h) = \text{diam}_\rho(S)$ , then we must have  $\rho(g_0 - f_k) = \text{diam}_\rho(S)$ , for  $k = 0, 1, \dots, n$ . This will contradict the maximality of  $S_0$ . Hence

$$\sup\{\rho(g - h) : g \in S\} = \rho(g_0 - h) < \text{diam}_\rho(S).$$

□

#### 4. MAIN RESULTS

**Theorem 4.1.** *Let  $\rho$  be a convex,  $\rho$  is a convex,  $\sigma$ -finite function modular satisfying the  $\Delta_2$ -type condition and  $C$  be a  $\rho$ -bounded,  $\rho$ -a.e. compact subset of  $L_\rho$ . Let  $T : C \rightarrow C$  be an asymptotically nonexpansive mapping. Let  $H$  be a convex subset of  $C$  such that:*

- (i) *if  $f \in H$  then  $\Omega_{\rho\text{-a.e.}}(f) \subset H$ ;*
- (ii) *for each  $f \in H$ , any subsequence  $\{T^{n_i}(f)\}_i$  of  $\{T^n(f)\}_n$ , has a  $\rho$ -convergent subsequence.*

*Then  $T$  has a fixed point.*

*Proof.* Consider the family  $\mathcal{F}$  of nonempty  $\rho$ -a.e. compact subset of  $H$  which satisfies property (3.1).  $\mathcal{F}$  is not empty since  $H \in \mathcal{F}$ . By the previous results,  $\mathcal{F}$  has a minimal element. Let  $K$  be a minimal element of  $\mathcal{F}$ . Assume that  $K$  has more than one point, i.e.  $\text{diam}_\rho(K) > 0$ . Let  $f \in K$ . Set

$$S = \Omega_{\|\cdot\|}(f) = \{g \in H; T^{n_i}(f) \|\cdot\|\text{-converges to } g \text{ for some } n_i \uparrow \infty\}.$$

It is easy to see that  $S \subset K$ . We claim that  $S = T(S)$ . Indeed, let  $g \in S$ . Then there exists a sequence  $\{T^{n_i}(f)\}_i$  which  $\|\cdot\|$ -converges to  $g$ . Since  $T$  is continuous, we have  $T^{n_i+1}(f) \xrightarrow{\|\cdot\|} T(g)$ . By definition of  $S$ , we get  $T(g) \in S$ , i.e.  $T(S) \subset S$ . Let us show the other inclusion, i.e.  $S \subset T(S)$ . Let  $g \in S$ . Again by definition of  $S$ , there exists a sequence  $\{T^{n_i}(f)\}_i$  which  $\|\cdot\|$ -converges to  $g$ . The sequence  $\{T^{n_i-1}(f)\}_i$  has a norm convergent subsequence, say  $\{T^{n_{\phi(i)}-1}(f)\}_i$ . Let  $h$  be its  $\|\cdot\|$ -limit. Since  $T$  is continuous, we get

$$T(h) = T(\lim_{i \rightarrow \infty} T^{n_{\phi(i)}-1}(f)) = \lim_{i \rightarrow \infty} T^{n_{\phi(i)}}(f) = g.$$

Hence  $g \in T(S)$ , i.e.  $S \subset T(S)$ . So our claim is proved, i.e.  $T(S) = S$ .

Next, notice that the assumption (ii) implies that  $S$  is norm compact. Lemma

(3.3) implies the existence of  $f_0 \in \overline{\text{conv}}(S) \subset K$  such that

$$\sup\{\rho(g - f_0) : g \in S\} < \text{diam}_\rho(S). \quad (A)$$

Let  $r = \sup\{\rho(g - f_0) : g \in S\}$ . Set

$$D = \{h \in K; \sup_{g \in S} \rho(g - h) \leq r\}.$$

Since  $f_0 \in D$  and  $\rho$  is convex,  $D$  is a nonempty convex subset of  $K$ . We claim that  $D = K$ . Indeed, let us first show that  $D$  is  $\rho$ -a.e. compact. By the assumption (ii), it is enough to show that  $D$  is  $\rho$ -a.e. closed. Let  $\{h_n\}_n$  be a sequence in  $D$  such that  $h_n \xrightarrow{\rho\text{-a.e.}} h \in L_\rho$ . Fix  $g \in S$ . Since  $g - h_n \xrightarrow{\rho\text{-a.e.}} g - h$ , Lemma (1.3) implies

$$\rho(g - h) \leq \liminf_{n \rightarrow \infty} \rho(g - h_n)$$

which yields

$$\rho(g - h) \leq \liminf_{n \rightarrow \infty} \left( \sup\{\rho(f - h_n) : f \in S\} \right) \leq r.$$

Hence  $\sup\{\rho(h - g) : g \in S\} \leq r$ , i.e.  $h \in D$ . Next we check that  $D$  satisfies property (3.1). Indeed, let  $f \in D$  and  $g \in \Omega_{\rho\text{-a.e.}}(f)$ . Then there exists a sequence  $\{T^{n_i}(f)\} \xrightarrow{\rho\text{-a.e.}} g$ . Using Lemma (1.3) we obtain

$$\rho(g - h) \leq \liminf_{n \rightarrow \infty} \rho(T^{n_i}(f) - h) \leq \limsup_{n \rightarrow \infty} \rho(T^n f - h)$$

for any  $h \in S$ . Since  $T(S) = S$ , there exists a sequence  $\{u_n\}_n$  in  $S$  such that  $h = T^n(u_n)$ , for any  $n \geq 1$ . Hence

$$\begin{aligned} \rho(g - h) &\leq \limsup_{n \rightarrow \infty} \rho(T^n f - T^n u_n) \leq \limsup_{n \rightarrow \infty} k_n \rho(f - u_n) \\ &\leq \limsup_{n \rightarrow \infty} \rho(f - u_n) \leq \sup\{\rho(f - u) : u \in S\} \leq r. \end{aligned}$$

So  $\sup\{\rho(g - h) : h \in S\} \leq r$  which gives  $g \in D$ . Thus  $D$  satisfies property (3.1) and by minimality of  $K$ , we obtain  $D = K$ . But

$$\text{diam}_\rho(D) \leq r < \text{diam}_\rho(S) \leq \text{diam}_\rho(K),$$

which is a contradiction. Therefore,  $K$  is reduced to one point. Property (3.1) will force this point to be a fixed point of  $T$ .  $\square$

Now we are ready to state and prove the main result of this work.

**Theorem 4.2.** *Let  $\rho$  be a convex,  $\rho$  is a convex,  $\sigma$ -finite function modular satisfying the  $\Delta_2$ -type condition and  $C$  be a convex  $\rho$ -bounded and  $\rho$ -a.e. compact subset of  $L_\rho$ . Let  $T : C \rightarrow C$  be  $\rho$ -asymptotically nonexpansive. Then  $T$  has a fixed point.*

*Proof.* Let  $\mathcal{F}$  be the family of nonempty convex subsets of  $C$  which satisfy the property (3.1).  $\mathcal{F}$  is not empty since  $C \in \mathcal{F}$ . By Zorn's lemma,  $\mathcal{F}$  has a minimal element. Let  $H$  be a minimal element of  $\mathcal{F}$ . Let us show that  $H$  satisfies the hypothesis of Theorem (4.1). It suffices to check that  $H$  satisfies property (ii). Let  $r$  be defined on  $H$  as in Lemma (3.2). If  $r = 0$  we have

$$\lim_{n \rightarrow \infty} T^n f = g$$

for any  $f, g \in H$ , which implies (ii). Otherwise, assume that  $r > 0$ . Let  $f \in H$  such that there exists a sequence  $\{T^{n_i} f\}_i$  which has no norm-convergent subsequence. Thus, there exists  $\varepsilon > 0$  and a subsequence  $\{T^{n(k)} f\}_k$  such that

$$\text{Sep}(\{T^{n(k)} f\}_k) = \inf\{\rho(T^{n(k)} f - T^{n(k')} f), k \neq k'\} \geq \varepsilon.$$

Since  $H$  is  $\rho$ -a.e. compact, there exists  $f_\infty \in H$  such that  $T^{n(k)} f \xrightarrow{\rho\text{-a.e.}} f_\infty \in H$  as  $k \rightarrow \infty$ . Without loss of generality, we may assume the existence of

$$\lim_{k \rightarrow \infty} \rho(T^{n(k)} f - f_\infty) = l.$$

Since  $\limsup_{n \rightarrow \infty} \rho(T^n f - f) = r$ , we choose  $\eta > 0$  such that  $\eta < \frac{\varepsilon}{2}$ , and an integer  $n_0 \geq 1$ , such that for all  $n \geq n_0$  we have

$$\rho(T^n f - f) < r + \eta.$$

Fix  $n \geq n_0$ . There exists  $k_0 \geq 1$  such that for all  $k \geq k_0$ , we have  $n(k) \geq n + n_0$  and

$$\begin{aligned} \rho(T^n f - T^{n(k)} f) &= \rho(T^n f - T^{n+(n(k)-n)} f) = \rho(T^n f - T^n(T^{n(k)-n} f)) \\ &\leq k_n \rho(f - T^{n(k)-n} f) < k_n(r + \eta). \end{aligned}$$

Note that if  $f_n \xrightarrow{\rho\text{-a.e.}} f$  and  $\text{Sep}\{f_n\}_n \geq \varepsilon$ , then by Lemma (1.3), we have

$$\varepsilon \leq \liminf_{m \rightarrow \infty} \liminf_{n \rightarrow \infty} \rho(f_n - f_m) \leq 2 \liminf_{n \rightarrow \infty} \rho(f_n - f).$$

Combined with Lemma (1.3), we get

$$\liminf_{n \rightarrow \infty} \rho(f_n) = \liminf_{n \rightarrow \infty} \rho(f_n - f) + \rho(f) \geq \frac{\varepsilon}{2} + \rho(f).$$

In particular, since  $\{T^{n(k)}f - T^n f\}_k$  is  $\rho$ -a.e. convergent to  $f_\infty - T^n f$  as  $k \rightarrow \infty$  and satisfies  $\text{Sep}(\{T^{n(k)}f - T^n f\}_k) \geq \varepsilon$ , we get

$$\rho(T^n f - f_\infty) \leq \liminf_{k \rightarrow \infty} \rho(T^{n(k)}f - T^n f) - \frac{\varepsilon}{2}.$$

Hence

$$\rho(f_\infty - T^n f) \leq r + \eta - \frac{\varepsilon}{2}$$

which implies

$$r = \limsup_{n \rightarrow \infty} \rho(f_\infty - T^n f) \leq r + \eta - \frac{\varepsilon}{2} < r.$$

This contradiction completes the proof of Theorem 4.2. □

Assume that  $L_\rho = L_p(\Omega, \mu)$  for a  $\sigma$ -finite measure  $\mu$ . If  $C$  is a convex, bounded and closed subset of  $L_p$  for  $1 < p < \infty$  and  $T : C \rightarrow C$  is asymptotically nonexpansive, it is known that  $C$  has a fixed point because  $L_p$  is uniformly convex. However the result does not hold for  $p = 1$  (even for nonexpansive mappings, see [1]). Since  $L_1$  is a modular space, Theorem (4.1) implies the existence of fixed point if  $p = 1$  when  $C$  is  $\rho$ -a.e. compact. Thus we can state.

**Corollary 4.1.** Let  $(\Omega, \mu)$  be as above,  $C \subset L_1(\Omega, \mu)$  a convex bounded set which is compact for the topology of the convergence local in measure and  $T : C \rightarrow C$  asymptotically nonexpansive. Then,  $T$  has a fixed point.

*Proof.* Under the above hypothesis  $\rho$ -a.e. compact sets and compact sets in the topology of convergence local in measure are identical. □

## REFERENCES

- [1] D.E. Alspach. *A fixed point free nonexpansive map*. Proc. Am. Math. Soc., (1981), **82**, 423-424.
- [2] S.C. Bose. *Weak convergence to the fixed point of an asymptotically nonexpansive map*. Proc. Am. Math. Soc., (1978), **68**, 305-308.
- [3] T. Dominguez Benavides, M.A. Khamsi, S. Samadi. *Asymptotically regular mappings in modular function spaces*. Preprint.
- [4] T.Dominguez Benavides, M.A. Khamsi, S. Samadi. *Uniformly Lipschitzian mappings in modular function spaces*. Preprint.
- [5] K. Goebel and W.A. Kirk. *A fixed point theorem for asymptotically nonexpansive mappings*. Proc. Am. Math. Soc., (1972), **35**, 171-174.
- [6] M.A. Khamsi. *Fixed point theory in modular function spaces*. Recent Advances on Metric Fixed Point Theory. Universidad de Sevilla, Sevilla, (1996), 31-58.
- [7] M.A. Khamsi, W.M. Kozłowski, S. Reich. *Fixed point theory in modular function spaces*. Nonlinear Anal., (1990), **14**, 935-953.
- [8] W.M. Kosłowski. *Modular function spaces*. Dekker: New York, Basel, (1988).
- [9] T.-H. Kim and H.-K. Xu. *Remarks on asymptotically nonexpansive mappings*. Nonlinear Anal., to appear.
- [10] C. Martínez Yañez. *A fixed point theorem on  $k$ -uniformly rotund spaces*. Nonlinear Anal., (1988), **13** 857-861.
- [11] G. Passty. *Construction of fixed points for asymptotically nonexpansive mappings*. Proc. Am. Math. Soc., (1982), **84**, 213-216.
- [12] H.-K. Xu.  *$k$ -Uniform rotundity and fixed points of mappings of asymptotically nonexpansive type*. To appear in Chinese.
- [13] H.-K. Xu. *Existence and convergence for fixed points of mappings of asymptotically nonexpansive type*. Nonlinear Anal., (1991) **16**(12), 1139-1146.
- [14] X.T. Yu and X. Dai. *A fixed point theorem of asymptotically nonexpansive mappings*. J. Math. (PRC), (1986), **6**, 255-262.

TOMAS DOMINGUEZ-BENAVIDES, DEPARTMENT OF MATHEMATICAL ANALYSIS, UNIVERSITY OF SEVILLE, P.O.BOX 1160. 41080. SEVILLE (SPAIN).

*E-mail address:* tomasd@cica.es

MOHAMED AMINE KHAMSI, DEPARTMENT OF MATHEMATICAL SCIENCE, THE UNIVERSITY OF TEXAS AT EL PASO, EL PASO, TX 79968, (U.S.A).

*E-mail address:* mohamed@math.utep.edu

SEDKI SAMADI, DEPARTMENT OF MATHEMATICAL ANALYSIS, UNIVERSITY OF SEVILLE,  
P.O.Box 1160. 41080. SEVILLE (SPAIN).

*E-mail address:* `sedki@cica.es`